

# Risk Aggregation under Dependence Uncertainty

Challenges in Theory and Practice

**Paul Embrechts**

RiskLab, Department of Mathematics, ETH Zurich

Senior SFI Professor

[www.math.ethz.ch/~embrechts/](http://www.math.ethz.ch/~embrechts/)

# Outline

- 1 Framework
- 2 VaR and ES Bounds
- 3 Asymptotic Equivalence
- 4 Challenges
- 5 References

# Fundamental problem in Finance/Insurance

- Risk factors:  $\mathbf{X} = (X_1, \dots, X_d)$
- Model assumption:  $X_i \sim F_i, F_i$  known,  $i = 1, \dots, d$
- A financial position  $\Psi(\mathbf{X})$
- A risk measure/pricing function:  $\rho(\Psi(\mathbf{X}))$

Calculate  $\rho(\Psi(\mathbf{X}))$

# Calculating $\rho(\Psi(\mathbf{X}))$

Example:

- $\Psi(\mathbf{X}) = \sum_{i=1}^d X_i$
- $\rho = \text{VaR}_p$  or  $\rho = \text{ES}_p$

Challenge:

- We need a *joint* model for the random vector  $\mathbf{X}$
- Joint models are hard to get by

We will focus on the above special choices of  $\Psi$  and  $\rho$ .

# VaR and ES

## VaR<sub>p</sub>(X)

For  $p \in (0, 1)$ ,

$$\text{VaR}_p(X) = F_X^{-1}(p) = \inf\{x \in \mathbb{R} : F_X(x) \geq p\}$$

## ES<sub>p</sub>(X)

For  $p \in (0, 1)$ ,

$$\text{ES}_p(X) = \frac{1}{1-p} \int_p^1 \text{VaR}_q(X) dq \stackrel{(F \text{ cont.})}{=} \mathbb{E}[X | X > \text{VaR}_p(X)]$$

# VaR and ES

A related quantity **Left-tail-ES**:

$LES_p(X)$

For  $p \in (0, 1)$ ,

$$LES_p(X) = \frac{1}{p} \int_0^p \text{VaR}_q(X) dq = -\text{ES}_{1-p}(-X)$$

# Fréchet problem

Denote

$$\mathcal{S}_d = \mathcal{S}_d(F_1, \dots, F_d) = \left\{ \sum_{i=1}^d X_i : X_i \sim F_i, i = 1, \dots, d \right\}$$

- Every element in  $\mathcal{S}_d$  is a possible risk position.
- Determination of  $\mathcal{S}_d$ : very challenging.
  - Think about  $\mathcal{S}_2(U[0, 1], U[0, 1])$  ... open question!

# Worst- and best-values of VaR and ES

The Fréchet (unconstrained) problems for  $\text{VaR}_p$

$$\overline{\text{VaR}}_p(S_d) = \sup\{\text{VaR}_p(S) : S \in \mathcal{S}_d(F_1, \dots, F_d)\},$$

$$\underline{\text{VaR}}_p(S_d) = \inf\{\text{VaR}_p(S) : S \in \mathcal{S}_d(F_1, \dots, F_d)\}.$$

Same notation for  $\text{ES}_p$  and  $\text{LES}_p$ .



# Worst- and best-values of VaR and ES

- ES is subadditive:

$$\overline{\text{ES}}_p(S_d) = \sum_{i=1}^d \text{ES}_p(X_i).$$

Similarly  $\underline{\text{LES}}_p(S_d) = \sum_{i=1}^d \underline{\text{LES}}_p(X_i)$ .

- $\overline{\text{VaR}}_p(S_d)$ ,  $\underline{\text{VaR}}_p(S_d)$  and  $\underline{\text{ES}}_p(S_d)$ : generally open questions

## Challenge for $\underline{\text{ES}}_p(S_d)$

To calculate  $\underline{\text{ES}}_p(S_d)$  one naturally seeks a **safest** risk in  $S_d$ .

# Mathematical difficulty

Common understanding of the **most dangerous** scenario:

- Comonotonicity - well accepted notion

Understanding concerning the **safest** scenario:

- $d = 2$ : counter-monotonicity
- $d \geq 3$ : question mark! (?!)
  - Calls for notions of extremal negative dependence.

# Mathematical difficulty

ES respects **convex order**: the natural order of risk preference.

## Convex order

We write  $X \leq_{cx} Y$  if  $\mathbb{E}[f(X)] \leq \mathbb{E}[f(Y)]$  for all convex functions  $f$  such that the two expectations exist.

## Finding $\underline{ES}_p(S_d)$

Search for a **smallest element** in  $\mathcal{S}_d$  with respect to convex order, if it exists.

# Mathematical difficulty

VaR does not respect convex order: more tricky

- Good news: the questions for  $\overline{\text{VaR}}_p(S_d)$ ,  $\underline{\text{VaR}}_p(S_d)$  and  $\underline{\text{ES}}_p(S_d)$  are mathematically similar.

## Finding $\overline{\text{VaR}}_p(S_d)$

Search for a **smallest element** in  $\mathcal{S}_d(\hat{F}_1, \dots, \hat{F}_d)$  with respect to convex order, where  $\hat{F}_i$  is the  $p$ -tail-conditional distribution of  $F_i$ .

- $\underline{\text{VaR}}_p(S_d)$  is symmetric to  $\overline{\text{VaR}}_p(S_d)$ .

# Summary of existing results

$d = 2$ :

- fully solved analytically

$d \geq 3$ :

- Homogeneous model ( $F_1 = \dots = F_d$ )
  - $\underline{ES}_p(S_d)$  solved analytically for decreasing densities, e.g. Pareto, Exponential
  - $\overline{VaR}_p(S_d)$  solved analytically for tail-decreasing densities, e.g. Pareto, Gamma, Log-normal
- Inhomogeneous model
  - Few analytical results: current research
- Numerical methods available: Rearrangement Algorithm

# VaR bounds

$d = 2$ , Makarov (1981) and Rüschendorf (1982)

For any  $p \in (0, 1)$ ,

$$\overline{\text{VaR}}_p(S_2) = \inf_{x \in [0, 1-p]} \{F_1^{-1}(p+x) + F_2^{-1}(1-x)\},$$

and

$$\underline{\text{VaR}}_p(S_2) = \sup_{x \in [0, p]} \{F_1^{-1}(x) + F_2^{-1}(p-x)\}.$$

- A **large outcome** is coupled with a **small outcome**.

# VaR bounds - homogeneous model

## Sharp VaR bounds (Wang, Peng and Yang, 2013)

Suppose that the density function of  $F$  is decreasing on  $[b, \infty)$  for some  $b \in \mathbb{R}$ . Then, for  $p \in [F(b), 1)$ , and  $X \stackrel{d}{\sim} F$ ,

$$\overline{\text{VaR}}_p(S_d) = d\mathbb{E}[X|X \in [F^{-1}(p + (d-1)c), F^{-1}(1-c)]],$$

where  $c$  is the smallest number in  $[0, \frac{1}{d}(1-p)]$  such that

$$\int_{p+(d-1)c}^{1-c} F^{-1}(t) dt \geq \frac{1-p-dc}{d} ((d-1)F^{-1}(p + (d-1)c) + F^{-1}(1-c)).$$

**Red part** clearly has an ES-type form.

- $c = 0$ :  $\overline{\text{VaR}}_p(S_d) = \overline{\text{ES}}_p(S_d)$ .

# VaR bounds - homogeneous model

## Sharp VaR bounds II

Suppose that the density function of  $F$  is decreasing on its support. Then for  $p \in (0, 1)$  and  $X \stackrel{d}{\sim} F$ ,

$$\underline{\text{VaR}}_p(S_d) = \max\{(d-1)F^{-1}(0) + F^{-1}(p), d\mathbb{E}[X|X \leq F^{-1}(p)]\}.$$

Red part has an LES form.



# ES bounds - homogeneous model

## Sharp ES bounds (Bernard, Jiang and Wang, 2014)

Suppose that the density function of  $F$  is decreasing on its support. Then for  $p \in (1 - dc, 1)$ ,  $q = (1 - p)/d$  and  $X \stackrel{d}{\sim} F$ ,

$$\begin{aligned}\underline{\text{ES}}_p(S_d) &= \frac{1}{q} \int_0^q \left( (d-1)F^{-1}((d-1)t) + F^{-1}(1-t) \right) dt, \\ &= (d-1)^2 \text{LES}_{(d-1)q}(X) + \text{ES}_{1-q}(X),\end{aligned}$$

where  $c$  is the smallest number in  $[0, \frac{1}{d}]$  such that

$$\int_{(d-1)c}^{1-c} F^{-1}(t) dt \geq \frac{1-dc}{d} \left( (d-1)F^{-1}((d-1)c) + F^{-1}(1-c) \right).$$

- One **large outcome** is coupled with  $d - 1$  **small outcomes**.

# Complete mixability

The homogeneous VaR and ES bounds are based on the notion of **complete mixability**:

## Complete mixability, Wang and Wang (2011)

A distribution function  $F$  on  $\mathbb{R}$  is called  $d$ -completely mixable ( $d$ -CM) if there exist  $d$  random variables  $X_1, \dots, X_d \sim F$  such that

$$\mathbb{P}(X_1 + \dots + X_d = dk) = 1,$$

for some  $k \in \mathbb{R}$ .

- Equivalently,  $\mathcal{S}_d(F, \dots, F)$  contains a constant.

# Complete mixability

- Some examples of  $d$ -CM distributions for all  $d \geq 2$ :  
Normal, Student t, Cauchy, Uniform.
- Most relevant result:  $F$  has a **monotone density on a finite interval** with a mean condition (depends on  $d$ ) is  $d$ -CM.
  - Examples: (truncated) Pareto, Gamma, Log-normal.
- Inhomogeneous version called **joint mixability**.
- A full characterization of these classes is at the moment is widely open.

# Numerical calculation

**Rearrangement Algorithm (RA):** Embrechts, Puccetti and Rüschendorf (2013).

- A fast numerical procedure
- Based on the CM-idea
- Discretization of relevant quantile regions
- $d$  possibly large
- Applicable to  $\overline{\text{VaR}}_p$ ,  $\underline{\text{VaR}}_p$  and  $\underline{\text{ES}}_p$

# Asymptotic equivalence

Consider the case  $d \rightarrow \infty$ . What would happen to  $\overline{\text{VaR}}_p(S_d)$ ?

- Clearly always  $\overline{\text{VaR}}_p(S_d) \leq \overline{\text{ES}}_p(S_d)$ .
- Recall that  $\overline{\text{VaR}}_p(S_d)$  has an ES-type part.

Under some weak conditions,

$$\lim_{d \rightarrow \infty} \frac{\overline{\text{ES}}_p(S_d)}{\overline{\text{VaR}}_p(S_d)} = 1.$$

This was shown first for homogeneous models and then extended to general inhomogeneous models.

# Asymptotic equivalence - homogeneous model

## Theorem 1

*In the homogeneous model,  $F_1 = F_2 = \dots = F$ , for  $p \in (0, 1)$  and  $X \sim F$ , we have that*

$$\lim_{d \rightarrow \infty} \frac{1}{d} \overline{\text{VaR}}_p(S_d) = \text{ES}_p(X).$$

- Similar limits hold for a large class of risk measures

# Asymptotic equivalence - worst-cases

## Theorem 2 (Embrechts, Wang and Wang, 2014)

Suppose the continuous distributions  $F_i$ ,  $i \in \mathbb{N}$  satisfy that for  $X_i \sim F_i$  and some  $p \in (0, 1)$ ,

(i)  $\mathbb{E}[|X_i - \mathbb{E}[X_i]|^k]$  is uniformly bounded for some  $k > 1$ ;

(ii)  $\liminf_{d \rightarrow \infty} \frac{1}{d} \sum_{i=1}^d \text{ES}_p(X_i) > 0$ .

Then as  $d \rightarrow \infty$ ,

$$\frac{\overline{\text{ES}}_p(S_d)}{\overline{\text{VaR}}_p(S_d)} = 1 + O(d^{1/k-1}).$$

# Asymptotic equivalence - best-cases

Similar results holds for  $\underline{\text{VaR}}_p$  and  $\underline{\text{ES}}_p$ : assume (i) and

$$(iii) \liminf_{d \rightarrow \infty} \frac{1}{d} \sum_{i=1}^d \text{LES}_p(X_i) > 0,$$

then

$$\lim_{d \rightarrow \infty} \frac{\underline{\text{VaR}}_p(S_d)}{\underline{\text{LES}}_p(S_d)} = 1,$$

$$\lim_{d \rightarrow \infty} \frac{\underline{\text{ES}}_p(S_d)}{\sum_{i=1}^d \mathbb{E}[X_i]} = 1,$$

and

$$\frac{\underline{\text{VaR}}_p(S_d)}{\underline{\text{ES}}_p(S_d)} \approx \frac{\sum_{i=1}^d \text{LES}_p(X_i)}{\sum_{i=1}^d \mathbb{E}[X_i]} \leq 1, \quad d \rightarrow \infty.$$



# Example: Pareto(2) risks

Bounds on VaR and ES for the sum of  $d$  Pareto(2) distributed rvs for  $p = 0.999$ ;  $\text{VaR}_p^+$  corresponds to the comonotonic case.

	$d = 8$	$d = 56$
$\underline{\text{VaR}}_p$	31	53
$\underline{\text{ES}}_p$	178	472
$\text{VaR}_p^+$	245	1715
$\overline{\text{VaR}}_p$	465	3454
$\overline{\text{ES}}_p$	498	3486
$\overline{\text{VaR}}_p / \text{VaR}_p^+$	1.898	2.014
$\overline{\text{ES}}_p / \overline{\text{VaR}}_p$	1.071	1.009

# Example: Pareto( $\theta$ ) risks

Bounds on the VaR and ES for the sum of  $d = 8$   
Pareto( $\theta$ )-distributed rvs for  $p = 0.999$ .

	$\theta = 1.5$	$\theta = 2$	$\theta = 3$	$\theta = 5$	$\theta = 10$
$\overline{\text{VaR}}_p$	1897	465	110	31.65	9.72
$\overline{\text{ES}}_p$	2392	498	112	31.81	9.73
$\overline{\text{ES}}_p / \overline{\text{VaR}}_p$	1.261	1.071	1.018	1.005	1.001

# Dependence-uncertainty spread

## Theorem 3 (Embrechts, Wang and Wang, 2014)

Take  $1 > q \geq p > 0$ . Suppose that the continuous distributions  $F_i$ ,  $i \in \mathbb{N}$ , satisfy (i) and (iii), and  $\limsup_{d \rightarrow \infty} \frac{\sum_{i=1}^d \mathbb{E}[X_i]}{\sum_{i=1}^d \text{ES}_p(X_i)} < 1$ , then

$$\liminf_{d \rightarrow \infty} \frac{\overline{\text{VaR}}_q(S_d) - \underline{\text{VaR}}_q(S_d)}{\overline{\text{ES}}_p(S_d) - \underline{\text{ES}}_p(S_d)} \geq 1.$$

- The **uncertainty spread** of VaR is generally bigger than that of ES.
- In recent Consultative Documents of the Basel Committee,  $\text{VaR}_{0.99}$  is compared with  $\text{ES}_{0.975}$ :  $p = 0.975$  and  $q = 0.99$ .

# Dependence-uncertainty spread

ES and VaR of  $S_d = X_1 + \dots + X_d$ , where

- $X_i \sim \text{Pareto}(2 + 0.1i)$ ,  $i = 1, \dots, 5$ ;
- $X_i \sim \text{Exp}(i - 5)$ ,  $i = 6, \dots, 10$ ;
- $X_i \sim \text{Log-Normal}(0, (0.1(i - 10))^2)$ ,  $i = 11, \dots, 20$ .




	$d = 5$			$d = 20$		
	best	worst	spread	best	worst	spread
$\text{ES}_{0.975}$	22.48	44.88	22.40	29.15	102.35	73.20
$\text{VaR}_{0.975}$	9.79	41.46	31.67	21.44	100.65	79.21
$\text{VaR}_{0.9875}$	12.06	56.21	44.16	22.12	126.63	104.51
$\text{VaR}_{0.99}$	12.96	62.01	49.05	22.29	136.30	114.01
$\frac{\text{ES}_{0.975}}{\text{VaR}_{0.975}}$		1.08			1.02	

# Challenges




Open mathematical questions:

- Characterization of complete and joint mixability
- Characterization of  $\mathcal{S}_d$
- Find  $\overline{\text{VaR}}_p$  under more general settings, especially in the inhomogeneous model
- Partial dependence information and realistic scenarios
- Marginal uncertainty and statistical estimation
- Many more ...



# References I

-  Bernard, C., X. Jiang, and R. Wang (2014). Risk aggregation with dependence uncertainty. *Insurance: Mathematics and Economics*, 54(1), 93–108.
-  Embrechts, P., G. Puccetti, and L. Rüschendorf (2013). Model uncertainty and VaR aggregation. *Journal of Banking and Finance*, 37(8), 2750–2764.
-  Embrechts, P., G. Puccetti, L. Rüschendorf, R. Wang, and A. Beleraj (2014). An academic response to Basel 3.5. *Risks*, 2(1), 25–48.

# References II

-  Embrechts, P., B. Wang, and R. Wang (2014). Aggregation-robustness and model uncertainty of regulatory risk measures. *Preprint*, ETH Zurich.
-  Makarov, G.D. (1981). Estimates for the distribution function of the sum of two random variables with given marginal distributions. *Theory of Probability and its Applications*, 26, 803–806.
-  Rüschenendorf, L. (1982). Random variables with maximum sums. *Advances in Applied Probability*, 14(3), 623–632.

# References III

-  Wang, B. and R. Wang (2011). The complete mixability and convex minimization problems with monotone marginal densities. *Journal of Multivariate Analysis* 102(10), 1344–1360.
-  Wang, R., L. Peng, and J. Yang (2013). Bounds for the sum of dependent risks and worst value-at-risk with monotone marginal densities. *Finance and Stochastics* 17(2), 395–417.



THANK YOU!